

**DRUG FACTS**

Active ingredient(s)

Purpose

- Use(s)
- temporarily reduces fever
  - temporarily relieves minor aches and pains due to:
    - headache
    - muscular aches
    - the common cold
    - minor pain of arthritis
    - backache
    - toothache
    - premenstrual and menstrual cramps

- Warnings
- Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 8 tablets in 24 hours, which is the maximum daily amount
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product

- Do not use
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
  - if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

have liver disease

Ask a doctor or pharmacist before use if you are

you are taking the blood thinning drug warfarin

When using this product

- Stop use and ask a doctor if
- pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of children

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Other information

adults and children 12 years and over	<ul style="list-style-type: none"><li>take 2 tablets every 4 to 6 hours while symptoms last</li><li>do not take more than 8 tablets in 24 hours</li><li>do not take for more than 10 days unless directed by a doctor</li></ul>
	do not use adult Extra Strength product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

store at 15° to 30°C (59° to 86°F)

Inactive ingredients

povidone, pregelatinized starch, stearic acid